

TruFi™ II Ultra 2x PCR Mix

Catalog No.	Pack Size and Concentration	Components and Volume
AZ-1920	100 x 50 µl reactions, 2x	TruFi™ II Ultra 2x PCR Mix - 2 x 1.25 mL
AZ-1921	500 x 50 µl reactions, 2x	TruFi™ II Ultra 2x PCR Mix - 10 x 1.25 mL
AZ-1922	1000 x 50 µl reactions, 2x	TruFi™ II Ultra 2x PCR Mix - 20 x 1.25 mL

Description

TruFi™ II Ultra 2x PCR Mix is an optimized 2x master-mix comprised of TruFi™ II DNA Polymerase, optimized reaction buffer, MgCl₂, and ultra-pure dNTPs. TruFi™ II Ultra 2x PCR Mix is ideal for fast cycling, as the polymerase exhibits a rapid elongation rate of 10 seconds per kb. The ready-to-use 2x mix is also highly efficient in long PCR with amplicons greater than 15 kb. With an error rate which is greater than 65x lower than Taq Polymerase, TruFi™ II Ultra 2x PCR Mix is recommended for applications which require extremely high fidelity such as cloning, NGS applications, and mutagenesis.

- **Ultra High-Fidelity:** provides fidelity that is >65x higher than Taq Polymerase and exhibits both 5'→ 3' DNA polymerase activity and 3'→ 5' proofreading exonuclease activity
- **Robust Amplification:** delivers greater yields with less optimization than other 2x master-mixes, even in amplification of GC-rich targets, DNA of sub-optimal purity, and long templates. TruFi™ II Ultra 2x PCR Mix has a high elongation rate of 10 sec/kb, which is 4x faster than *Taq*

Storage

TruFi™ II Ultra 2x PCR Mix is shipped on blue ice and should be stored at –20°C upon receipt. Excessive freeze/thawing should be avoided.

Important Guidelines

TruFi™ II Ultra 2x PCR Mix: The 2x mix is comprised of TruFi™ II DNA Polymerase, optimized reaction buffer, 3.0 mM MgCl₂, and ultra-pure dNTPs. The composition has been developed to deliver maximum efficiency, sensitivity and success with most amplicons. Completely thaw 2x mix and thoroughly mix by inversion for proper resuspension of precipitate. In some cases, a higher concentration of MgCl₂ or the addition of 1M -2M betaine can improve performance.

Template: For complex genomic DNA, we suggest the use of 5 ng – 500 ng per reaction; For cDNA or plasmid DNA, please use < 100 ng per reaction.

Primers: Primers should have a predicted melting temperature of around 60°C, using default Primer 3 settings (<http://frodo.wi.mit.edu/primer3/>). The final primer concentration in the reaction should be between 0.2 µM and 0.6 µM.

Annealing: We recommend performing a temperature gradient to determine the optimal annealing temperature. Alternatively, 58°C can be used as a starting point. For optimization, increase in 2°C increments. Due to the high salt concentration in the 2x mix, the optimal annealing temperature may be higher than with traditional PCR buffers.

Extension: Optimal extension is achieved at 72°C. The optimal extension time is dependent on amplicon length and complexity. 10 to 30 seconds per kilobase (kb) is recommended for amplification from eukaryotic genomic DNA or cDNA. For simple targets such as plasmid DNA and short targets less than 2 kb, 10 seconds per kb is typically sufficient.

Reaction setup – Allow TruFi™ II Ultra 2x PCR Mix to thaw and mix contents thoroughly by inversion.

1. Prepare a PCR mixture on ice based on following table, adding template DNA to individual tubes containing PCR mixture:

Component	50 µl Reaction	Final Concentration/Notes
TruFi™ II Ultra 2x PCR Mix	25 µl	1x
Forward Primer (10 µM)	2.0 µl	400 nM
Reverse Primer (10 µM)	2.0 µl	400 nM
Template DNA	<100 ng cDNA, <500 ng genomic DNA	variable
PCR-grade water	Up to 50 µl final volume	

* For alternative total reaction volumes (eg. 25 µl), scale all components proportionally to maintain final concentrations.

2. PCR cycling:

Cycles	Temperature & Time	Notes
1	98°C, 2 minutes	Initial Denaturation
25 - 40	98°C, 15 seconds 58°C to 66°C, 15 seconds 72°C, 10 to 30 seconds per kb	Denaturation Annealing* (determined by user) Extension

* We suggest performing a temperature gradient to determine the optimal annealing temperature. See Important Guidelines.

Quality Control

TruFi™ II Ultra 2x PCR Mix is tested extensively for robust activity, processivity, efficiency, heat activation, sensitivity, absence of nuclease contamination and absence of nucleic acid contamination. TruFi™ II Ultra 2x PCR Mix is manufactured under a comprehensive quality management system, following ISO 9001:2015 standards.

Limitations of Use

This product is intended for research purposes only and is not intended for any animal or human therapeutic use.

Technical Support

For Trouble-shooting and Technical Guidance, please contact us at tech@azuragenomics.com and provide PCR reaction conditions, cycling parameters, amplicon size, and screen grabs (gel images) if possible.

Azura Genomics guarantees the performance of all products in the manner described in our product literature. The purchaser must determine the suitability of the product for its particular use. Azura Genomics shall not be liable for any direct, indirect, consequential or incidental damages arising from the use, the result of use or the liability to use this product.